

RESEARCH ARTICLE

Open Access

Association of knowledge, preventive counseling and personal health behaviors on physical activity and consumption of fruits or vegetables in community health workers

Alex A Florindo^{1*}, Ross C Brownson², Gregore I Mielke³, Grace AO Gomes⁴, Diana C Parra⁵, Fernando V Siqueira⁶, Felipe Lobelo⁷, Eduardo J Simoes⁸, Luiz R Ramos⁹, Mário M Bracco¹⁰ and Pedro C Hallal⁶

Abstract

Background: There is evidence that if a health professional is active and has a healthy diet, he/she is more likely to advise patients about the benefits of physical activity and healthy eating. The aims of this study were to: (1) describe the personal physical activity, consumption of fruits and vegetables behaviors and nutritional status of community health workers; (2) evaluate the association between knowledge, delivery of preventive counseling and personal behaviors among community health workers.

Methods: This was a cross-sectional study conducted in a nationally sample of health professionals working in primary health care settings in Brazil in 2011. This survey was part of the second phase of the Guide for Useful Interventions for Activity in Brazil and Latin America project, and data were collected through telephone interviews of 269 community health workers from the Unified Health Care system of Brazil. We applied questionnaires about personal reported behaviors, knowledge and preventive counseling in physical activity and consumption of fruits and vegetables. We calculated the prevalence and associations between the variables with logistic regression.

Results: The proportion of community health workers that practiced 150 minutes per week of physical activity in leisure time or transportation was high (64.9%). Half of community health workers were overweight and only 26.2% reported consuming five portions/day of fruits or vegetables. Most community health workers reported counseling about physical activity for more than six months (59.7%), and most were not knowledgeable of the fruits and vegetables and physical activity recommendations. Meeting the fruits and vegetables recommendations was associated with correct knowledge (OR = 4.5; CI95% 1.03;19.7), with reporting 150 minutes or more of physical activity per week (OR = 2.0; CI95% 1.03;3.7) and with reporting physical activity in leisure time (OR = 2.0; CI95% 1.05;3.6). Regular physical activity counseling was associated with reporting 10–149 minutes per week (OR = 3.8; CI95% 1.1;13.3) and with more than 150 minutes of physical activity per week (OR = 4.9; CI95% 1.5;16.5).

Conclusion: Actions to promote physical activity and healthy eating and to improve knowledge among community health workers within the health care system of Brazil could have a potential positive influence on delivery of preventive counseling to patients on this topic.

Keywords: Physical activity, Consumption of fruits and vegetables, Community health workers, Primary health care settings, Knowledge, Preventive counseling, Personal behavior

* Correspondence: aflorind@usp.br

¹School of Arts, Sciences and Humanities, University of São Paulo, Rua Arlindo Bettio, 1.000, CEP: 03828-000 São Paulo, SP, Brazil

Full list of author information is available at the end of the article

Background

Physical inactivity, unhealthy diets, and obesity are among the top ten risk factors for morbidity and mortality worldwide [1]. This situation is particularly concerning in low and middle-income countries. In Brazil, for example, more than half of adult population is overweight, and 60% do not practice any kind of physical activity in leisure time, nor they do report regular consumption of fruits and vegetables [2,3]. A wide range of research and interventions have been implemented aimed at changing this scenario [4,5], but low and middle-income countries still have few policies implemented for promoting physical activity and fruits and vegetables intake [6]. In a recent study published by our research group, it was found that 39.8% and 72% of the primary care units in Brazil, develop any physical activity or healthy eating program, respectively [7].

The implementation of interventions in primary health care settings is a possible strategy for addressing the burden of obesity-related diseases [8]. Many of these interventions have focused on the training of health professionals, mainly physicians and nurses, to provide counseling for physical activity and healthy eating. Such interventions have shown positive results at improving health behaviors of patients, [9,10] and are facilitated by improving knowledge about the benefits of health promotion and improving counseling skills among health professionals in the primary care setting [10]. In addition, studies have shown that if the health professional is active and has a healthy diet, he/she is more likely to advise patients about the benefits of physical activity and healthy eating [11-15].

Most studies on this topic have been conducted with physicians and nurses [9-13]. In the last 10 years, community health workers have been shown to also play a key role in health promotion and disease prevention [16-25]. In the Brazilian context, community health workers living and working in the same neighborhood are typically responsible for visiting families in their homes in order to provide more personal guidance and counseling for disease prevention and health promotion, orientation about health services, and strengthen the linkage between health professionals teams and families. The health family team includes a physician, a nurse, auxiliary-nurses and a community health worker [26]. The Brazilian Ministry of Health has a guide that includes proposals of actions for the community health workers in Brazilian municipalities. This document contains orientations of counseling in healthy food, physical activity and weight control for health [27]. However, there is a lack of information about counseling and behaviors in physical activity, healthy food and obesity/weight control among community health workers, as well as associations between these factors. Therefore, the

aims of this study were to: (1) describe the behaviors of physical activity, consumption of fruits and vegetables, and nutritional status of community health workers; (2) evaluate the associations between recommendations and knowledge of healthy behaviors (physical activity, fruits and vegetables consumption and body mass index) with personal healthy behaviors (fruits and vegetables consumption, body mass index and physical activity) among community health workers.

Methods

We conducted a cross-sectional study with a nationally-representative sample of primary health care units in Brazil in 2011. This survey was part of second phase of the GUIA project in Brazil (Guide for Useful Interventions for Activity in Brazil and Latin America). We drew a random sample in three stages: (1) 1,600 primary health care units were selected randomly of a total 42,486 units in Brazil; (2) within the 1,600 sampled primary health care units, we listed three types of professionals: physicians, nurses and community health workers; (3) in 1/3 of the primary health care units we tried to interview the physicians, in 1/3 the nurses and in 1/3 the community health workers. A detail description of the sampling strategy of primary health care units and health professionals is available elsewhere [28]. We used random digit dialing survey to interview community health workers between January and July of 2011. We were able to interview one person in 1,279 primary health care units (79.9% of the approached ones). We interviewed 269 community health workers since only in one third of the units, the community health worker was the professional chosen. In addition, several of the units did not have a community health worker, thus the response rate of 50.5% should be interpreted with caution. The questionnaire was applied by telephone and included sociodemographic variables (age, gender, education, skin color, marital status), smoking, self-rated health, self-reported behaviors such as physical activity practice and consumption of fruits and vegetables, nutritional status, knowledge and professional conduct in physical activity, knowledge in consumption of fruits and vegetables and body mass index for classification of nutritional status.

Variable description

Physical activity

We assessed the domains of leisure and transportation physical activity through self-report using the long version of the International Physical Activity Questionnaire (IPAQ). A validation study of this version of IPAQ administered by both face-to-face and telephone interviews of adults in Brazil is published [29]. We classified self-reported physical activity based on the recommendation

of the World Health Organization [30]: a cutoff of 150 minutes per week of moderate-to-vigorous physical activity was applied, summing time spent active in the leisure time and transportation domains: 1) Physically active: practice of 150 minutes or more per week; 2) Insufficiently active: practice of 10 to 149 minutes per week; 3) Inactive: < 10 minutes per week. We also categorized practice of 150 minutes or more per week (yes or no); practice of leisure time physical activity (yes or no) and practice of transportation physical activity (yes or no).

Consumption of fruits and vegetables

We used two questions to classify self-reported consumption of fruits and vegetables: (1) How many portions of fruits the community health workers consumed per day on average; (2) How many portions of vegetables the community health workers consumed per day on average. If the interviewee had doubts about the meaning of one portion, the interviewer used examples (speaking of an apple or banana for example). Natural juices and vegetable soups were also included. We classified community health workers consuming at least five portions per day of fruits or vegetables as having a healthy diet according to the World Health Organization recommendation or otherwise [31].

Body mass index

We calculated body mass index (BMI) based on self-reported weight and height. We categorized BMI using the World Health Organization standards [31]: (1) normal weight: $BMI \leq 24.9 \text{ kg/m}^2$; (2) overweight: $BMI \geq 25.0 \text{ kg/m}^2$; (3) obesity: $BMI \geq 30.0 \text{ kg/m}^2$.

Knowledge about physical activity, consumption of fruits and vegetables and nutritional status

We assessed knowledge about physical activity, consumption of fruits and vegetables, and nutritional status based on the recommendations from the World Health Organization [31], according to the following criteria:

- 1) Physical activity: We used two open questions, community health workers could select the answer that they considered to be correct: (1) Days per week of practice of moderate-intensity physical activity needed to achieve health benefits. The answer considered as adequate was a minimum of five days; (2) Time per day of practice of moderate-intensity physical activity needed to achieve health benefits. The answer considered as adequate was a minimum of 30 minutes per day. We classified knowledge of physical activity as a practice of at least 30 minutes per day of moderate activities in at least five times per week. This questionnaire has been used in one

intervention study with community health workers in Brazil [32].

- 2) Consumption of fruits and vegetables: We used one closed question of multiple choice about the minimum of portions of fruits and vegetables that an adult should consume per day with. The question had seven options. The correct choice was a minimum of five portions per day of fruits or vegetables.
- 3) Weight status: We applied one multiple choice closed question regarding cutoff points for overweight and obesity. The question had five options. The correct choice was $BMI \geq 25.0 \text{ kg/m}^2$ for overweight and $BMI \geq 30.0 \text{ kg/m}^2$ for obesity.

Professional advice/counseling for promoting physical activity

We used one question about community health workers' habit to give advice about physical activity to patients, with six possible answers: (1) Does not advise and does not plan to start; (2) Does not advise, but does intend to start; (3) Does advise, but not regularly; (4) Does advise, but started recently; (5) Advises regularly, for more than six months; (6) Used to advise, but stopped doing so. We categorized professional practice for community health workers that positive response to option 5 (advises regularly, for more than six months). This questionnaire was already used in an intervention study among community health workers in Brazil [32].

Data analysis

We measured the prevalence of community health workers' self-reported physical activity, consumption of fruits and vegetables and nutritional status; their preventive counseling for physical activity; and knowledge about recommendation for physical activity, knowledge about recommendation for consumption of fruits and vegetables and the knowledge about cutoff for classification of overweight and obesity. Initially we calculated the bivariate association between all variables using chi-square test: 1) leisure time physical activity (yes or no); 2) transportation physical activity (yes or no); 3) practice of at least 150 minutes of physical activity per week (yes or no); 4) practice of physical activity in three levels (practice of 150 minutes or more per week; practice of 10 to 149 minutes per week; inactive or < 10 minutes per week); 5) consumption of at least five portions per day of fruits or vegetables (yes or no); 6) $BMI (\geq 25.0 \text{ kg/m}^2 \text{ or } \leq 24.9 \text{ kg/m}^2)$; 7) physical activity counseling for more than six months (yes or no); 8) knowledge about physical activity recommendations (know or not know); 9) knowledge about the recommendations of fruits and vegetables consumption (know or not know); 10) knowledge about cutoff of BMI for classification overweight and

obesity (know or not know). All variables that had significant association for $p < 0.05$ were calculated logistic regression models and odds ratio, unadjusted and adjusted by gender, age and education. The selection of dependent variables for regression models was done based in two criteria: 1) variables that were more associated between all of them; 2) variables that were dichotomized. All analysis were performed using SPSS software, version 15.0.

Ethical issues

The study was approved by the Ethics Committee of the Federal University of Pelotas (process number 16154).

Results

The respondents in our sample of community health workers were mainly women, ages 49 years and older, had less than 11 years of education, had non-white skin color, were married or living with a partner, had good or very good health, and were not smokers (Table 1). Most

Table 1 Sociodemographic and general health characteristics of community health workers, Brazil, 2011

Variables	Community health workers %
Gender	
Men	10.8
Women	89.2
Education (years of study)	
Up to 8 years	7.8
9 to 11 years	79.6
12 years or more	12.6
Age	
18 to 29 years old	28.3
30 to 49 years old	61.3
50 years or more	10.4
Skin Color*	
White	39.0
Non-white	61.0
Marital status	
Single	24.2
Married/Living with partners	66.9
Widowed/Divorced/Separated	8.9
Self-report of health	
Bad or very bad	20.1
Good or very good	79.9
Smoking	
Yes	7.4
No	92.6

*2 missing.

community health workers practiced leisure time or transportation physical activity and achieved physical activity recommendations for health (Figure 1), but did not consume five portions per day of fruits or vegetables. Half of them were overweight (Figure 2).


Most community health workers were not aware of the recommendations of physical activity and consumption of fruits and vegetables for health as well as BMI cut off values for the classification of overweight and obesity. Over half of community health workers reported advising physical activity to people of their covered area (Figure 3).

Five significant associations with $p < 0.05$ were identified from bivariate analysis: (1) consumption of at least five portions per day of fruits or vegetables (yes or no) was associated with knowledge about the recommendations of fruits and vegetables consumption (know or not know) ($p = 0.018$); (2) leisure time physical activity (yes or no) was associated with consumption of at least five portions per day of fruits or vegetables (yes or no) ($p = 0.023$); (3) practice of at least 150 minutes of physical activity per week (yes or no) was associated with consumption of at least five portions per day of fruits or vegetables (yes or no) ($p = 0.038$); (4) practice of physical activity in three levels (practice of 150 minutes or more per week; practice of 10 to 149 minutes per week; inactive or < 10 minutes per week) was associated with consumption of at least five portions per day of fruits or vegetables (yes or no) ($p = 0.018$); (5) practice of physical activity in three levels (practice of 150 minutes or more per week; practice of 10 to 149 minutes per week; inactive or < 10 minutes per week) was associated with physical activity counseling for more than six months (yes or no) ($p = 0.011$).

We opted to put the consumption of at least five portions per day of fruits or vegetables (yes or no) and physical activity counseling for more than six months (yes or no) as dependent variables in logistic regression models, producing unadjusted and adjusted odds ratios (Table 2). Knowledge about recommendations for consumption of fruits and vegetables, just as practice of leisure time physical activity and the practice of regular physical activity for health were associated with higher odds ratios for consumption of five portions per day of fruits or vegetables. The practice of some physical activity or according to recommendations for health was associated with higher odds ratio for physical activity counseling. While three-level physical activity was associated with consumption of five portions of fruits or vegetables in bivariate analysis (p -trend), this association disappeared after adjustment in logistic models.

Discussion


This study showed that healthy behaviors reported, such as physical activity and consumption of fruits and




vegetables, were associated with knowledge and preventive counselling among community health workers in Brazil. To our knowledge, this is the first study that showed that physical activity practice was associated with physical activity counseling among community health workers.

Health promotion interventions that include community health workers are important because these professionals have the potential to influence the behavior of patients from having a closer contact with them within the primary health care system [16-24]. Although most community health workers practice physical activity at recommended levels, most do not consume fruits and vegetables in accordance with recommendations, and half were overweight. In comparison with the Brazilian adult population, community health workers practice more physical activity, but the consumption of fruits and vegetables as well as overweight and obesity prevalence were similar [2,3]. Interventions to improve health promotion of the community health workers can result in

more effective professional actions for patients and the whole population who are covered by them. A controlled study conducted with health professionals in Israel showed that an intervention for physicians and dietitians resulted in significant reduction of waist circumference and improved knowledge for counseling about weight loss (physicians) and healthy lifestyle (dietitians) [33]. In the same study, nurses and health promotion specialists did not change physical activity levels or healthy eating behaviors or showed reduction in waist circumference, which shows the complexity of changing behaviors [33]. A systematic review conducted with physicians and nurses showed that the association between weight status and management practices for weight control in patients is weak [34]. In our study with community health workers, we also found no association between knowledge about the correct classification of overweight or obesity and BMI. Four years of intervention conducted by Frank et al. to increase physical activity, consumption of fruits and vegetables and cessation of smoking and





alcohol consumption among medical students, resulted in better preparation for counseling for physical activity and healthy eating among these students [15].

This study showed a high prevalence of community health workers who did not know the recommendations of physical activity and consumption of fruits and vegetables for health. Half these community health workers were overweight and most of them do not consumed five portions of fruits and vegetables per day. Therefore, interventions to improve knowledge and health behaviors related to physical activity and healthy eating are needed among community health workers.

Other studies have shown that physicians and nurses of primary health care settings in Brazil [28] as well as general practitioners, practice nurses and health visitors in Scotland did not know the recommendations of physical activity for health [35]. We have evidence that interventions can improve knowledge and professional conduct among community health workers [32,36] and they are similar among other health professionals [15,37]. Community health workers are more usually women of lower socioeconomic status. These professionals and others who are in stressful jobs (prone to burnout and poorer health) are sometimes overlooked in interventions to promote health [38,39]. Therefore, interventions to improve lifestyle of community health workers are needed.

The results of this study showed that although the prevalence of physical activity counseling was high, community health workers had little knowledge about a recommendation or global strategy for physical activity promotion and healthy diet of the World Health Organization. The high prevalence of physical activity counseling is possibly influenced by physical activity promotion policies implemented by the Brazilian Ministry

of Health during the last years and encouraged in cities and health care services for health promotion [40,41]. But the little knowledge about physical activity indicates that community health workers need a more organized and comprehensive training. We believe that there is not a standard for community health workers to advise in physical activity in primary health care system in Brazil. For example, the practice guide of community health workers [27] needs to be improved with better information about physical activity promotion and the Brazilian Ministry of Health could encourage the municipalities in training these professionals. Some models of training for physical activity promotion by community health workers has been studied by researchers in Brazil and could be applied [32,42]. This could contribute in actions to fight chronic diseases for reduce physical inactivity [43].

The study has some limitations. The cross-sectional data does not allow us to determine temporality and thus move toward establishing. The methods for evaluation of knowledge were not validated against "gold standards" (e.g., for preventive counseling). A study published recently among community health workers used a similar questionnaire to evaluate the knowledge about physical activity showing that this instrument was adequate for evaluating changes in physical activity promotion [32]. Another limitation is the high non-response rate of 50.5% and the missing information about community health workers in primary health care units. Not all primary health care units had community health workers, because professionals in some units use a different model (the health family model). This fact may have introduced bias toward a non-representative sample. In addition, these professionals may have had time constraints for response and lack of telephone lines in

Table 2 Relationships of knowledge about healthy behaviors and preventive counseling for physical activity self-reported healthy behaviors among community health workers, Brazil, 2011

	Reported consuming five portions per day of fruits or vegetables	
	OR (CI95%) ¹	OR (CI95%) ²
<i>Correct knowledge of daily fruits and vegetables to consumption recommendation (five portions per day)</i>		
No	1	1
Yes	5.0 (1.2 – 21.4)	4.5 (1.03 – 19.7)
<i>Reported physical activity (minutes per week)</i>		
<150 minutes	1	1
150 minutes or more	2.0 (1.03 – 3.7)	2.0 (1.03 – 3.7)
<i>Leisure time physical activity reported</i>		
No	1	1
Yes	2.0 (1.1 – 3.7)	2.0 (1.05 – 3.6)
<i>Reported physical activity (minutes per week)</i>		
Physical inactivity (<10 minutes)	1	1
Practice of 10 to 149 minutes	4.0 (0.5 – 32.3)	4.0 (0.5 – 33.0)
Practice of 150 minutes or more	6.5 (0.8 – 50.6)	6.5 (0.83 – 51.2)
Counseling on regular physical activity (at least six months) (yes)		
OR (CI95%)¹ OR (CI95%)²		
<i>Reported physical activity (minutes per week)</i>		
Physical inactivity (<10 minutes)	1	1
Practice of 10 to 149 minutes	3.4 (1.0 – 11.8)	3.8 (1.1 – 13.3)
Practice of 150 minutes or more	4.7 (1.4 – 15.5)	4.9 (1.5 – 16.5)

OR (odds ratio); CI (confidence interval) ¹Unadjusted; ²Adjusted for gender, age and education.

some primary health care units. However, the sample of this study is very similar to other samples of community health workers in Brazil in relation to age, education and gender [44–46], minimizing the possibility of selection bias.

Conclusions

This study showed that most community health workers in Brazil practice physical activity and advise their patients to physical activity, but do not know the recommended amount of physical activity needed to obtain health benefits. Most community health workers did not know recommendations of the consumption of fruits and vegetables, did not consume at least five portions per day of fruits and vegetables, and half are overweight. We found an association between knowledge, preventive counselling and healthy behaviors

reported of physical activity and consumption of fruits and vegetables. Therefore, actions to health promotion are needed for community health workers. On the other hand, it seems promising that community health workers can disseminate physical activity and healthy eating counseling into the communities where they serve, with appropriate training.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AAF and PCH contributed to the idea this study. AAF, ROB, GIM, FVS, GAOG and PCH contributed for data analysis, interpretation and drafted the manuscript. DCP, LRR, FL and MMB contributed to drafting and critically revising the manuscript. All authors have read and approved the final manuscript.

Acknowledgments

This study was funded through the Centers for Disease Control and Prevention's Prevention Research Centers Program special interest project U48/DP001903 (Applying Evidence-Physical Activity Recommendations in Brazil).

Author details

¹School of Arts, Sciences and Humanities, University of São Paulo, Rua Arlindo Bettio, 1.000, CEP: 03828-000 São Paulo, SP, Brazil. ²Prevention Research Center in St. Louis, Brown School, Division of Public Health Sciences and Alvin J. Siteman Cancer Center - Washington University in St. Louis, Missouri, USA. ³Postgraduate Program in Epidemiology, Federal University of Pelotas, Pelotas, RS, Brazil. ⁴Department of Gerontology, Federal University of São Carlos, São Carlos, SP, Brazil. ⁵Program in Physical Therapy Surgery (Prevention and Control), School of Medicine, Washington University in St. Louis, St. Louis, MO, USA. ⁶School of Physical Education, Federal University of Pelotas, Pelotas, RS, Brazil. ⁷Hubert Department of Global Health, Emory University - Rollins School of Public Health, Atlanta, USA. ⁸Department of Health Management and Informatics, School of Medicine, University of Missouri, Missouri, USA. ⁹Department of Preventive Medicine, Federal University of São Paulo, São Paulo, SP, Brazil. ¹⁰Hospital Israelita Albert Einstein; Hospital Municipal Dr. Moisés Deutsch - M'Boi Mirim, São Paulo, SP, Brazil.

Received: 10 November 2014 Accepted: 16 March 2015

Published online: 09 April 2015

References

- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380(9859):2224–60.
- Florindo AA, Hallal PC, Moura EC, Malta DC. Practice of physical activities and associated factors in adults, Brazil, 2006. *Rev Saude Publica*. 2009;43 Suppl 2:65–73.
- Ministério da Saúde do Brasil. Vigilância de fatores de risco e proteção para doenças crônicas por inquérito telefônico, VIGITEL, BRASIL, 2012. Brasília, DF: Secretaria de Vigilância em Saúde. Departamento de Vigilância de Doenças e Agravos não transmissíveis e Promoção da Saúde; 2013. in portuguese.
- Jaime PC, da Silva AC, Gentil PC, Claro RM, Monteiro CA. Brazilian obesity prevention and control initiatives. *Obes Rev*. 2013;14 Suppl 2:88–95.
- Kohl 3rd HW, Craig CL, Lambert EV, Inoue S, Alkandari JR, Leetongin G, et al. The pandemic of physical inactivity: global action for public health. *Lancet*. 2012;380(9838):294–305.
- Lachat C, Otchere S, Roberfroid D, Abdulai A, Seret FM, Milesevic J, et al. Diet and physical activity for the prevention of noncommunicable diseases in low- and middle-income countries: a systematic policy review. *PLoS Med*. 2013;10(6):e1001465.
- Ramos LR, Malta DC, Gomes GA, Bracco MM, Florindo AA, Mielke GI, et al. Prevalence of health promotion programs in primary health care units in Brazil. *Rev Saude Publica*. 2014;48(5):837–44.

8. Orrow G, Kinmonth AL, Sanderson S, Sutton S. Effectiveness of physical activity promotion based in primary care: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2012;344:e1389.
9. ter Bogt NC, Milder IE, Bemelmans WJ, Beltman FW, Broer J, Smit AJ, et al. Changes in lifestyle habits after counselling by nurse practitioners: 1-year results of the Groningen Overweight and Lifestyle study. *Public Health Nutr*. 2011;14(6):995–1000.
10. Grandes G, Sanchez A, Montoya I, Ortega Sanchez-Pinilla R, Torcal J, Group P. Two-year longitudinal analysis of a cluster randomized trial of physical activity promotion by general practitioners. *PLoS One*. 2011;6(3):e18363.
11. Wells KB, Lewis CE, Leake B, Ware Jr JE. Do physicians preach what they practice? A study of physicians' health habits and counseling practices. *JAMA*. 1984;252(20):2846–8.
12. Frank E, Wright EH, Serdula MK, Elon LK, Baldwin G. Personal and professional nutrition-related practices of US female physicians. *Am J Clin Nutr*. 2002;75(2):326–32.
13. Frank E, Bhat Schelbert K, Elon L. Exercise counseling and personal exercise habits of US women physicians. *J Am Med Womens Assoc*. 2003;58(3):178–84.
14. Huijig JM, Gebhardt WA, Verheijden MW, van der Zouwe N, de Vries JD, Middelkoop BJ, et al. Factors influencing primary health care professionals' physical activity promotion behaviors: a systematic review. *Int J Behav Med*. 2015;22(1):32–50.
15. Frank E, Elon L, Hertzberg V. A quantitative assessment of a 4-year intervention that improved patient counseling through improving medical student health. *MedGenMed*. 2007;9(2):58.
16. Balcazar HG, Wise S, Redelfs A, Rosenthal EL, de Heer HD, Burgos X, et al. Perceptions of Community Health Workers (CHWs/PS) in the U.S.-Mexico Border HEART CVD study. *Int J Environ Res Public Health*. 2014;11(2):1873–84.
17. Rothschild SK, Martin MA, Swider SM, Tumalian Lynas CM, Janssen I, Avery EF, et al. Mexican American trial of community health workers: a randomized controlled trial of a community health worker intervention for Mexican Americans with type 2 diabetes mellitus. *Am J Public Health*. 2014;114(8):1540–8.
18. Jennings A, Barnes S, Okereke U, Welch A. Successful weight management and health behaviour change using a health trainer model. *Perspect Public Health*. 2013;133(4):221–6.
19. Coleman KJ, Farrell MA, Rocha DA, Hayashi T, Hernandez M, Wolf J, et al. Readiness to be physically active and self-reported physical activity in low-income Latinas, California WISEWOMAN, 2006–2007. *Prev Chronic Dis*. 2012;9:e87.
20. Spinner JR, Alvarado M. Salud Para Su Carozon—a Latino promotora-led cardiovascular health education program. *Fam Community Health*. 2012;35(2):111–9.
21. Staten LK, Cutshaw CA, Davidson C, Reinschmidt K, Stewart R, Roe DJ. Effectiveness of the Pasos Adelante chronic disease prevention and control program in a US-Mexico border community, 2005–2008. *Prev Chronic Dis*. 2012;9:E08.
22. Babamoto KS, Sey KA, Camilleri AJ, Karlan VJ, Catalasan J, Morisky DE. Improving diabetes care and health measures among hispanics using community health workers: results from a randomized controlled trial. *Health Educ Behav*. 2009;36(1):113–26.
23. Gibbons MC, Tyus NC. Systematic review of U.S.-based randomized controlled trials using community health workers. *Prog Community Health Partnersh*. 2007;1(4):371–81.
24. Staten LK, Gregory-Mercado KY, Ranger-Moore J, Will JC, Giuliano AR, Ford ES, et al. Provider counseling, health education, and community health workers: the Arizona WISEWOMAN project. *J Womens Health*. 2004;13(5):547–56.
25. Johnson CM, Sharkey JR, Dean WR, St John JA, Castillo M. Promotoras as research partners to engage health disparity communities. *J Acad Nutr Diet*. 2013;113(5):638–42.
26. Ministério da Saúde do Brasil. Política Nacional de Atenção Básica. Brasília, DF: Secretaria de Atenção à Saúde. Departamento de Atenção Básica; 2012. in portuguese.
27. Ministério da Saúde do Brasil. Guia Prático do Agente Comunitário de Saúde. Brasília, DF: Secretaria de Atenção à Saúde. Departamento de Atenção Básica; 2009. in portuguese.
28. Florindo AA, Mielke GI, Gomes GA, Ramos LR, Bracco MM, Parra DC, et al. Physical activity counseling in primary health care in Brazil: a national study on prevalence and associated factors. *BMC Public Health*. 2013;13:794.
29. Hallal PC, Simoes E, Reichert FF, Azevedo MR, Ramos LR, Pratt M, et al. Validity and reliability of the telephone-administered international physical activity questionnaire in Brazil. *J Phys Act Health*. 2010;7(3):402–9.
30. World Health Organization. Global recommendations on physical activity for health. Geneva; 2010.
31. World Health Organization. Diet, nutrition and the prevention of chronic diseases: report of a joint WHO/FAO expert consultation. Geneva: World Technical Report Series 916; 2003.
32. Florindo AA, Costa EF, Sa TH, Santos TI, Velardi M, Andrade DR. Physical activity promotion in primary health care in Brazil: a counseling model applied to community health workers. *J Phys Act Health*. 2014;11:1531–9.
33. Shahar DR, Henkin Y, Rozen GS, Adler D, Levy O, Safra C, et al. A controlled intervention study of changing health-providers' attitudes toward personal lifestyle habits and health-promotion skills. *Nutrition*. 2009;25(5):532–9.
34. Zhu DQ, Norman IJ, While AE. The relationship between doctors' and nurses' own weight status and their weight management practices: a systematic review. *Obes Rev*. 2011;12(6):459–69.
35. Douglas F, Torrance N, van Teijlingen E, Meloni S, Kerr A. Primary care staff's views and experiences related to routinely advising patients about physical activity. A questionnaire survey. *BMC public health*. 2006;6:138.
36. Sranachareonpong K, Hanning RM. Diabetes prevention education program for community health care workers in Thailand. *J Community Health*. 2012;37(3):610–8.
37. Spencer EH, Frank E, Elon LK, Hertzberg VS, Serdula MK, Galuska DA. Predictors of nutrition counseling behaviors and attitudes in US medical students. *Am J Clin Nutr*. 2006;84(3):655–62.
38. Ge C, Fu J, Chang Y, Wang L. Factors associated with job satisfaction among Chinese community health workers: a cross-sectional study. *BMC Public Health*. 2011;11:884.
39. Lopes DMQ, Beck CLC, Prestes FC, Weiller TH, Colomé JS, Silva GMD. Community health agents and their experiences of pleasure and distress at work: a qualitative study. *Rev Esc Enferm USP*. 2012;46(3):633–40.
40. Malta DC, BarbosadaSilva J. Policies to promote physical activity in Brazil. *Lancet*. 2012;380(9838):195–6.
41. Knuth AG, Malta DC, Cruz DK, Castro AM, Fagundes J, Sardinha LM, et al. Description of the countrywide physical activity network coordinated by the Brazilian Ministry of Health: 2005–2008. *J Phys Act Health*. 2010;7 Suppl 2:S253–8.
42. Gomes MA, Duarte MFS. Efetividade de uma intervenção de atividade física em adultos atendidos pela estratégia saúde da família: programa ação e saúde Floripa - Brasil. *Rev Bras Ativ Fis Saúde*. 2008;13(1):44–56. in portuguese.
43. Ministério da Saúde do Brasil. Plano de Ações Estratégicas para o Enfrentamento das Doenças Crônicas não Transmissíveis no Brasil, 2011–2022. Brasília, DF: Secretaria de Vigilância em Saúde. Departamento de Análise de Situação de Saúde. Coordenação Geral de Doenças e Agravos Não Transmissíveis; 2011. in portuguese.
44. Nascimento CMB. Análise do cumprimento das práticas dos agentes comunitários de saúde em municípios da Região Metropolitana do Recife. (Dissertação de Mestrado). Fundação Oswaldo Cruz. Centro de Pesquisa Aggeu Magalhães, Recife, PE. Mestrado em Saúde Pública; 2008 in portuguese.
45. Ferraz L, Aerts DRGC. O Cotidiano de trabalho do Agente Comunitário de Saúde no PSF em Porto Alegre. *Ciêns Saúde Colet*. 2005;10(2):347–55. in portuguese.
46. Lino MM, Lanzoni MM, Albuquerque GL, Schweitzer MC. Perfil socioeconômico, demográfico e de trabalho dos Agentes Comunitários de Saúde. *Cogitare Enferm*. 2012;17(1):57–64. in portuguese.